

## 베르노니아 시네레아 지상부의 화학 성분 및 항염증 활성

윤의중\*† · 장영기\*\*

\*극지연구소 극지생명과학연구부, \*\*하와이주립대학교 약학대학

### Chemical Constituents from the Aerial Parts of *Vernonia cinerea* L. and Their Anti-Inflammatory Activity

Ui Joung Youn\*† and Leng Chee Chang\*\*

\*Division of Life Sciences, Korea Polar Research Institute, KIOST, Incheon 21990, Korea.

\*\*Department of Pharmaceutical Sciences, Daniel K. Inouye College of Pharmacy, University of Hawaii, Hilo, HI 96720, USA.

#### ABSTRACT

**Background:** Previous phytochemical studies of whole *Vernonia cinerea* L. plants have identified sesquiterpene lactones, sterols, and triterpenes, which possess anticancer, antifeedant, and antimalarial activities. However, there are no reports of other types of bioactive metabolites. Therefore, the present study aimed to identify phenolic compounds with anti-inflammatory activity in the aerial parts of the plant.

**Methods and Results:** Compounds were isolated from the aerial parts of *V. cinerea* using a silica and C-18 gel columns and semi-preparative HPLC instrument, and the structures of the compounds were determined using one- and two- dimension nuclear magnetic resonance spectroscopy and mass spectroscopy. The chloroform soluble extracts and isolated compounds were evaluated for their anti-inflammatory potential based on their ability to inhibit nitric oxide production and TNF- $\alpha$  induced NF- $\kappa$ B activity.

**Conclusions:** Phytochemical study of the aerial parts of *V. cinerea* led to the isolation of six phenolic compounds. Compound 1 was a major metabolite, and to the best of our knowledge, compounds 2 - 6 were isolated from *V. cinerea* for the first time. Among the isolates, compounds 1 and 3 exhibited TNF- $\alpha$ -induced NF- $\kappa$ B activity with IC<sub>50</sub> values of 7.5 and 11.5 M, respectively, and the inhibitory activity of phenyl propanoid compound 3 on TNF- $\alpha$ -induced NF- $\kappa$ B was evaluated for the first time.

**Key Words:** *Vernonia cinerea* L., Anti-Inflammatory Activity, Asteraceae, Phenolic Compounds

#### INTRODUCTION

NF- $\kappa$ B is a transcription factor that is associated with cell apoptosis, differentiation, and migration. Upon activation, it may promote cell proliferation and prevent cell death through anti-apoptotic factors (Baldwin, 2001). Inhibition of NF- $\kappa$ B signalling has a potential application for either the treatment or prevention of cancer.

Nitric oxide (NO) is an inorganic gaseous molecule that is synthesized by the oxidation of L-arginine catalyzed by

nitric oxide synthase (NOS) and is involved in a number of physiological and pathological processes in mammals (Moncada *et al.*, 1991). In the NOS family, iNOS is expressed in a variety of cells including macrophages, endothelial cells, and smooth muscle cells in response to pro-inflammatory stimuli such as IL-1 $\beta$ , TNF- $\alpha$ , and LPS. NO plays an important role in the regulation of many physiological functions, such as host defence, neurotoxicity, and vasodilation (Anggård, 1994). However, excess production of NO has been implicated in immunological and

†Corresponding author: (Phone) +82-32-760-5562 (E-mail) ujyou@kopri.re.kr

Received 2016 October 5 / 1st Revised 2016 November 12 / 2nd Revised 2016 December 6 / 3rd Revised 2016 December 12 / Accepted 2016 December 12

This is an open access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

inflammatory diseases including septic shock, rheumatoid arthritis, graft rejection, and diabetes (Anggård, 1994). Therefore, inhibition of NO increase is apparently an important therapeutic consideration in the development of anti-inflammatory and cancer agents.

The leaves, roots, and seeds of some species of *Vernonia* (Asteraceae) have been reported to have medicinal properties (Misra *et al.*, 1984a). *Vernonia cinerea* L. is an annual herb that grows in south Asia. The plant is called 'Yaa Saam Wan' in Thailand. It is used as a tonic, stomachic, and astringent and also known to treat consumption, asthma, and bronchitis (Kone and Kande, 2012). Aqueous ethanolic extracts showed anti-ranikhet virus and anti-cancer activities (Beeran *et al.*, 2014). The roots are used as an anthelmintic and diuretic (Dastur, 1977). The flowers are used to treat conjunctivitis, fever, and rheumatism (Kirtikar and Basu, 1975). There have been phytochemical reports on the diverse compounds from this species, including sesquiterpene lactones, sterols, and triterpenes (Kuo *et al.*, 2003; Chea *et al.*, 2006; Misra *et al.*, 1984a, b). Some of these compounds have been shown to have anticancer (Kuo *et al.*, 2003; Pratheeshkumar and Kuttan, 2011), antimalarial (Chea *et al.*, 2006), and antifeedant activity (Tandon *et al.*, 1998).

In a continuing research for anti-inflammatory agents, the chloroform (CHCl<sub>3</sub>)-soluble extract of the aerial parts of *V. cinerea* exhibited inhibitory effects on the tumor necrosis factor alpha (TNF- $\alpha$ )-induced NF- $\kappa$ B activity and lipopolysaccharide (LPS)-induced nitric oxide (NO) production using murine macrophage RAW 264.7 cells. Although previous phytochemical studies on the whole plant of *V. cinerea* have reported anticancer, antimalarial, and antifeedant activity, there are no reports on other type of secondary metabolites with their anti-inflammatory activity. In particular, phenolic compounds, such as phenolic acids, flavonoids, and tannins are a large group of phytochemicals, existing ubiquitously in plants as secondary metabolites, which exhibit a wide range of biological and physiological functions, such as anti-allergenic, anti-inflammatory, anti-microbial and antioxidant activities (Balasundram *et al.*, 2006; Manach *et al.*, 2005; Middleton *et al.*, 2000).

Accordingly, this research aims to find phenolic compounds with their anti-inflammatory activity, which have never been reported from the plant source. Here, we report the isolation and structure elucidation of secondary

metabolites, as well as the inhibition on TNF- $\alpha$ -induced NF- $\kappa$ B activity and nitric oxide (NO) production in LPS-stimulated RAW 264.7 cells.

## MATERIALS AND METHODS

### 1. General experiments

UV spectra were recorded on a Shimadzu PharmaSpec-1700 UV-visible spectrophotometer (Shimadzu, Kyoto, Japan). IR spectra were measured on a Bruker Tensor-27 spectrophotometer (Bruker, Billerica, MA, USA). 1D and 2D NMR spectra were recorded on a Bruker AVANCE (400 MHz) spectrometer (Bruker, Billerica, MA, USA). Low- and high-resolution mass spectrometer analyses were performed with a BioTOF II ESI mass spectrometer (Bruker, Billerica, MA, USA). Thin-layer chromatography (TLC) was performed on silica gel 60 F<sub>254</sub> (0.25 mm, Merck, Darmstadt, Germany). Silica gel (230-400 mesh, Merck, Darmstadt, Germany) and C-18 (YMC GEL ODS-A, 12 nm, S-150  $\mu$ m, YMC Co., Ltd., Kyoto, Japan) were used for column chromatography. Semi-preparative HPLC (Beckman Coulter Inc., Brea, CA, USA) was conducted on a Beckman Coulter Gold-168 system equipped with a photodiode array detector using an Alltech reversed-phase Econosil C-18 column (10  $\mu$ m, 10  $\times$  250 mm, Alltech Inc., Nicholasville, KY, USA) with a flow rate of 2 mL/min.

### 2. Plant material

The aerial parts of *Vernonia cinerea* L. (Asteraceae) were collected from the Lampang Herb Conservation Club, Lampang Province, Thailand, in May 2011 and identified by comparison with the voucher specimen at the Forest Herbarium, Bangkok, Thailand. A voucher specimen (no. Vcw 002) was deposited at the Natural Product Chemistry Laboratory, College of Pharmacy, University of Hawaii at Hilo.

### 3. Extraction and isolation

The air dried aerial parts of *V. cinerea* (1 kg) were extracted with MeOH (2 times  $\times$  2  $\ell$ ) at room temperature. The solvent was concentrated in vacuo to yield a MeOH extract (80 g), which was then suspended in distilled water (0.5  $\ell$ ) and fractionated with chloroform (CHCl<sub>3</sub>, 2  $\times$  1  $\ell$ ), ethyl acetate (EtOAc, 2  $\times$  1  $\ell$ ), and *n*-butanol (BuOH, 2  $\times$  1  $\ell$ ), successively.

The  $\text{CHCl}_3$  extracts (12 g) were subjected to silica gel column chromatography (CC;  $\varnothing$  10 cm; 230 - 400 mesh, 0.5 kg) using a gradient solvent system of hexane-ethyl acetate (100:0 to 30:70), to afford 57 fractions (C1-C57). The fractions (1 g) combined from C54 to C57 were subjected to silica gel CC ( $\varnothing$  8 cm; 230 - 400 mesh, 80 g), with hexane-ethyl acetate (100:0 to 1:1) as the solvent system, yielding seven subfractions (C54S1 to C54S7). Subfraction C54S4 (0.5 g) was chromatographed on a sephadex LH-20 gel (50 g) column and eluted with  $\text{H}_2\text{O}$ -MeOH (100:0 to 50:50), to afford five subfractions (C54S4L1 to C54S4L5), and compound 3 (1.5 mg, 0.000019% recovery from the extract) was recrystallized from subfraction C54S4L2. Subfraction C54S4L3 (0.1 g) was subjected to preparative. HPLC (MeOH- $\text{H}_2\text{O}$  = 40:60 to 100:0) to yield 6 (5 mg,  $t_R$ : 98 min, 0.00005%), 5 (2 mg,  $t_R$ : 105 min, 0.000025%), 4 (1 mg,  $t_R$ : 108 min, 0.000012%), and 2 (4 mg,  $t_R$ : 115 min, 0.00005%).

Quercetin 3-*O*- $\beta$ -D-glucopyranoside, 1 (30 mg, 0.00038%) was recrystallized in a solvent mixture, ( $\text{CHCl}_3$ :MeOH, 1:1) from the ethyl acetate portion.

**1) Quercetin 3-*O*- $\beta$ -D-glucopyranoside** - Yellow amorphous powder, UV (MeOH)  $\lambda_{\text{max}}$  (log  $\epsilon$ ) 279 (4.0), 330 (3.7) nm; IR  $\nu_{\text{max}}$  (KBr) 3325, 1655  $\text{cm}^{-1}$ ; ESI-MS  $m/z$  465  $[\text{M} + \text{H}]^+$ ;  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  7.48 (1H, dd,  $J=8.0, 2.0$  Hz, H-6'), 7.46 (1H, d,  $J=8.0$  Hz, H-5'), 6.94 (1H, d,  $J=2.0$  Hz, H-2'), 6.82 (1H, d,  $J=2.0$  Hz, H-8), 6.48 (1H, d,  $J=2.0$  Hz, H-6), 5.12 (1H, d,  $J=7.2$  Hz, H-1"), 3.75 (1H, d,  $J=10.4$  Hz, H-6a"), 3.53 (1H, d,  $J=10.4$  Hz, H-6b"), 3.20-3.53 (4H, m, H-2" to H-5"), 12.96 (1H, s, OH-5);  $^{13}\text{C}$  NMR (100 MHz,  $\text{DMSO}-d_6$ )  $\delta$  182.3 (C-4), 164.9 (C-7), 163.4 (C-5), 161.6 (C-9), 157.4 (C-2), 150.4 (C-4'), 146.2 (C-3'), 121.8 (C-1'), 119.6 (C-6'), 116.6 (C-5'), 114.0 (C-2'), 105.8 (C-10), 103.6 (C-1"), 100.3 (C-6), 95.1 (C-8), 77.6 (C-3"), 76.8 (C-5"), 73.5 (C-2"), 70.0 (C-4"), 61.0 (C-6").

**2) (*E*)-4-(3,4-Dimethoxyphenyl)but-3-en-1-ol** - Yellow amorphous powder, UV (MeOH)  $\lambda_{\text{max}}$  (log  $\epsilon$ ) 280 (4.1) nm; ESI-MS  $m/z$  209  $[\text{M} + \text{H}]^+$ ;  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  6.94 (1H, d,  $J=2.0$  Hz, H-2'), 6.91 (1H, dd,  $J=7.4, 2.0$  Hz, H-6'), 6.82 (1H, d,  $J=7.4$  Hz, H-5'), 6.44 (1H, d,  $J=16.4$  Hz, H-4), 6.09 (1H, dt,  $J=16.4, 6.8$  Hz, H-3), 3.77 (1H, t,  $J=6.6$  Hz, H-1), 2.50 (1H, q,  $J=6.6$  Hz, H-2), 3.92 (3H, s, 3'- $\text{OCH}_3$ ), 3.89 (3H, s, 4'- $\text{OCH}_3$ );  $^{13}\text{C}$  NMR (100 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  149.1 (C-4'), 148.8 (C-3'),

132.1 (C-4), 130.5 (C-1'), 124.2 (C-6'), 124.2 (C-3), 111.3 (C-5'), 108.8 (C-2'), 62.0 (C-1), 36.1 (C-2), 55.7 (4'- $\text{OCH}_3$ ), 55.6 (3'- $\text{OCH}_3$ ).

**3) 3-Hydroxy-1-(4-hydroxy-3-methoxyphenyl)-propan-1-one** - White amorphous powder, UV (MeOH)  $\lambda_{\text{max}}$  (log  $\epsilon$ ) 254 (3.6) nm; ESI-MS  $m/z$  197  $[\text{M} + \text{H}]^+$ ;  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  7.57 (1H, dd,  $J=8.4, 2.0$  Hz, H-6'), 6.98 (1H, d,  $J=8.4$  Hz, H-5'), 6.93 (1H, d,  $J=2.0$  Hz, H-2'), 4.04 (2H, t,  $J=5.2$  Hz, H-3), 3.98 (3H, s, 3'- $\text{OMe}$ ), 3.20 (2H, t,  $J=5.2$  Hz, H-2);  $^{13}\text{C}$  NMR (100 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  199.0 (C-1), 151.0 (C-4'), 147.4 (C-3'), 131.2 (C-1'), 124.4 (C-6'), 115.8 (C-2'), 112.1 (C-5'), 59.0 (C-9'), 42.0 (C-8'), 56.7 (3'- $\text{OMe}$ ).

**4) 1*H*-Indole-3-carbaldehyde** - White amorphous powder, ESI-MS  $m/z$  146  $[\text{M} + \text{H}]^+$ ;  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  9.90 (1H, s, CHO), 8.85 (1H, dd,  $J=6.8, 1.2$  Hz, H-4), 8.17 (1H, s, H-2), 7.48 (1H, dd,  $J=6.8, 1.2$  Hz, H-7), 7.28 (2H, m, H-5/H-6);  $^{13}\text{C}$  NMR (100 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  185.5 (CHO), 138.0 (C-2), 137.0 (C-1a), 124.3 (C-6), 124.0 (C-3a), 122.5 (C-5), 121.5 (C-4), 117.0 (C-3), 112.2 (C-7).

**5) *trans*-Cinnamic acid** - Yellow amorphous powder, UV (MeOH)  $\lambda_{\text{max}}$  (log  $\epsilon$ ) 280 (4.0) nm; ESI-MS  $m/z$  149  $[\text{M} + \text{H}]^+$ ;  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  7.72 (1H, d,  $J=16.0$  Hz, H-3), 7.60 (2H, m, H-2'/H-6'), 7.42 (3H, m, H-3'/H-4'/H-5'), 6.50 (1H, d,  $J=16.0$  Hz, H-2);  $^{13}\text{C}$  NMR (100 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  172.0 (C-1), 147.9 (C-3), 115.3 (C-3), 134.2 (C-1'), 129.3 (C-3'/C-5'), 125.5 (C-2'/C-6'), 127.4 (C-4').

**6) Uracil** - White amorphous powder, ESI-MS  $m/z$  113  $[\text{M} + \text{H}]^+$ ;  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  5.45 (1H, d,  $J=7.6$  Hz, H-5), 7.39 (1H, d,  $J=7.6$  Hz, H-6);  $^{13}\text{C}$  NMR (100 MHz,  $\text{DMSO}-d_6$ )  $\delta$  164.7 (C-4), 151.7 (C-2), 142.5 (C-6), 100.4 (C-5).

#### 4. Tumor necrosis factor- $\alpha$ (TNF- $\alpha$ ) activated nuclear factor-kappa B (NF- $\kappa$ B) assay

Human embryonic kidney cells 293 Panomics (Fremont, CA, USA) were employed for monitoring changes occurring along the NF- $\kappa$ B pathway (Kondratyuk *et al.*, 2012). Stable constructed cells were seeded into 96-well plates at  $20 \times 10^3$  cells per well. Cells were maintained in Dulbecco's modified Eagle's medium (DMEM) (Invitrogen Co., Carlsbad, CA, USA), supplemented with 10% fetal bovine serum (FBS), 100 units/ml penicillin, 100  $\mu\text{g}/\text{ml}$

streptomycin, and 2 mM l-glutamine. After 48 h incubation, the medium was replaced and the cells were treated with various concentrations of test substances. TNF- $\alpha$  (Human, Recombinant, *E. coli*, Calbiochem, San Diego, CA, USA) was used as an activator at a concentration of 2 ng/ml (0.14 nM). The plate was incubated for 6 h. Spent medium was discarded and the cells were washed once with PBS. Cells were lysed using 50  $\mu$ l (for 96-well plate) reporter lysis buffer from Promega, by incubating for 5 min on a shaker, and stored at  $-80^{\circ}\text{C}$ . The luciferase assay was performed using the Luc assay system from Promega (Promega Co., Madison, WI, USA). The gene product, luciferase enzyme, reacts with luciferase substrate, emitting light which was detected using a luminometer (LUMIstar Galaxy BMG, Ortenberg, Germany). Data for NF- $\kappa$ B constructs are expressed as  $\text{IC}_{50}$  values (i.e. concentration required to inhibit TNF-activated NF- $\kappa$ B activity by 50%). As a positive control, two known NF- $\kappa$ B inhibitors were used: TPCK,  $\text{IC}_{50}=3.8 \mu\text{M}$  and BAY-11,  $\text{IC}_{50}=2.0 \mu\text{M}$ .

### 5. Inhibition of nitric oxide (NO) production in lipopolysaccharide (LPS)-activated murine macrophage RAW 264.7 cells (iNOp) assay

The level of nitrite, the stable end product of NO, was estimated as described previously (Park *et al.*, 2011). Briefly, RAW 264.7 cells were seeded and incubated in 96-well culture plates at  $37^{\circ}\text{C}$ , 5%  $\text{CO}_2$  in a humidified air for 24 h. The cultured medium was replaced with phenol red-free medium containing various concentrations of compounds for 15 min prior to 1 mg/ml of LPS exposure for 20 h. The amount of nitrite in the culture media was measured by using Griess reagent. Under the same experimental conditions, SRB assays were performed to evaluate the cytotoxic effect of compounds toward RAW 264.7 cells. L-N<sup>G</sup>-monomethyl arginine citrate (l-NMMA), as a positive control of this assay showed an  $\text{IC}_{50}$  value of 25.1  $\mu\text{M}$ .

### 6. Statistical analysis

All data were expressed as means  $\pm$  SD. Statistical analysis was performed using SPSS software (version 18.0, SPSS Inc., Chicago, IL, USA). Significant differences between the groups were determined using the analysis of variance (ANOVA) followed by Duncan Post Hoc Test (DPHT).  $p < 0.05$  were considered as statistical significance.

## RESULTS AND DISCUSSION

### 1. Structure elucidation of compounds 1 - 6

Repeated chromatography of the aerial parts of *V. cinerea* on silica gel and YMC-pack RP-C<sub>18</sub> columns led to the isolation of six known compounds (1 - 6) (Fig. 1).

Compound 1 was obtained as a light yellow amorphous powder and its molecular weight was evaluated as  $m/z$  465  $[\text{M} + \text{H}]^+$  by the positive ESI-MS spectrometry. The UV spectrum showed absorption maxima at 330 and 279 nm indicating a conjugated double bond and a carbonyl group, respectively. The IR spectrum displayed characteristic absorption bands at 3,325 and 1,655  $\text{cm}^{-1}$ , corresponding to the hydroxy group (s) and conjugated C=O group (s), respectively. The  $^{13}\text{C}$  NMR spectrum revealed 15 carbon signals, along with six glucose unit carbons at [ $\delta$  103.6 (C-1''), 77.6 (C-3''), 76.8 (C-5''), 73.5 (C-2''), 70.0 (C-4''), 61.0 (C-6'')], suggested that 1 is a glycosylated flavone skeleton (Han *et al.*, 2004). The  $^1\text{H}$  NMR spectrum showed a downfield shifted OH proton at  $\delta_{\text{H}}$  12.96 (1H, s, OH-5) due to the hydrogen bond with a carbonyl group (C-4), and two aromatic doublet protons at  $\delta_{\text{H}}$  6.82 (1H, d,  $J=2.0$  Hz, H-8) and 6.48 (1H, d,  $J=2.0$  Hz, H-6), suggesting a di-substituted A-ring. In addition, the  $^1\text{H}$  NMR spectrum displayed ABX type aromatic system protons at [ $\delta_{\text{H}}$  7.48 (1H, dd,  $J=8.0, 2.0$  Hz, H-6'), 7.46 (1H, d,  $J=8.0$  Hz, H-5'), and 6.94 (1H, d,  $J=2.0$  Hz, H-2')] in B-ring and glucose unit protons at [ $\delta_{\text{H}}$  5.12 (1H,




Fig. 1. Structures of compounds 1 - 6 isolated from *V. cinerea*.

d,  $J=7.2$  Hz, H-1''), 3.75 (1H, d,  $J=10.4$  Hz, H-6''a), 3.53 (1H, d,  $J=10.4$  Hz, H-6''b), 3.20-3.53 (4H, m, H-2'' to H-5''). The heteronuclear multiple bond correlation (HMBC) correlation of the anomeric proton (H-1'') with C-3 indicated the position of glucose unit at C-3. Accordingly, compound 1 was assigned as quercetin 3-*O*- $\beta$ -D-glucopyranoside (1), by comparison of its physicochemical data with literature values (Choi *et al.*, 2000).

Compound 2 was obtained as a yellow amorphous powder. The ESI-MS spectrum of 2 gave a molecular ion  $[M+H]^+$  at  $m/z$  209. The NMR and heteronuclear single-quantum correlation (HSQC) spectra of 2 displayed ABX-type aromatic protons at [ $\delta_H$  6.94 (1H, d,  $J=2.0$  Hz, H-2'), 6.91 (1H, dd,  $J=7.4, 2.0$  Hz, H-6'), and 6.82 (1H, d,  $J=7.4$  Hz, H-5')] with six aromatic carbons at [ $\delta_C$  149.1 (C-4'), 148.8 (C-3'), 130.5 (C-1'), 124.2 (C-6'), 111.3 (C-5'), and 108.8 (C-2')], a *trans*-olefinic group at  $\delta_H$  6.44 (1H, d,  $J=16.4$  Hz/ $\delta_C$  132.1 (C-4) and at 6.09 (1H, dt,  $J=16.4, 6.8$  Hz/ $\delta_C$  124.2 (C-3), and two methoxy groups at  $\delta_H$  3.92/ $\delta_C$  55.6 (3'-OCH<sub>3</sub>) and at  $\delta_H$  3.89/ $\delta_C$  55.7 (4'-OCH<sub>3</sub>). The position of two methoxy signals at ( $\delta_H$  3.92 and 3.89) was confirmed at C-3' and C-4', respectively, through the HMBC analysis (Fig. 2). The remaining methylene at  $\delta_H$  2.50 (1H, q,  $J=6.6$  Hz)/ $\delta_C$  36.1 (C-2) and an oxygen bearing methylene group at  $\delta_H$  3.77 (1H, t,  $J=6.6$  Hz)/ $\delta_C$  62.0 (C-1) were confirmed as a part of the *trans*-olefinic group, on the basis of the HMBC correlations from H-1 to C-3 and from H-2 to C-4. In addition, three-bond HMBC correlations of the olefinic proton (H-4) with C-2'/C-6' and of H-2'/H-6' with C-4 were shown as in Fig. 2, indicating a but-3-en-1-ol group was attached at C-1' of the phenyl group. Accordingly, the structure of 2 was identified as (*E*)-4-(3,4-dimethoxyphenyl)but-3-en-1-ol (Masuda and Jitoe, 1995).

Compound 4 was obtained as a white amorphous powder. The MS spectrum showed a molecular ion peak at  $m/z$  146  $[M+H]^+$ . The NMR and HSQC spectra of 4 displayed signals for a tri-substituted olefinic group at  $\delta_H$  8.17 (1H, s)/ $\delta_C$  (138.0, C-2) and  $\delta_C$  (117.0, C-3), an aromatic quaternary carbon at  $\delta_C$  124.0 (C-3a), a downfield shifted aromatic quaternary at  $\delta_C$  137.0 (C-1a) due to a nitrogen atom attachment, and four protonated aromatic signals at [ $\delta_H$  8.85 (1H, dd,  $J=6.8, 1.2$  Hz)/ $\delta_C$  121.5 (C-4), 8.17 (1H, s)/ $\delta_C$  138.0 (C-2), 7.48 (1H, dd,  $J=6.8, 1.2$  Hz)/ $\delta_C$  112.2 (C-7),  $\delta_H$  7.28 (1H, m)/ $\delta_C$  122.5




Fig. 2. Key  $^1\text{H}$ - $^{13}\text{C}$  HMBC correlations for 2 and 4.

(C-5), and  $\delta_H$  7.28 (1H, m)/ $\delta_C$  124.3 (C-6)], indicated the presence of an indole skeleton (Wang *et al.*, 2013). In addition, the NMR spectra revealed an aldehyde group signal at  $\delta_H$  9.90 (1H, s)/ $\delta_C$  185.5 (CHO). The HMBC spectrum displayed two- to three-bonds correlations from H-2 to aldehyde carbon (CHO)/C-3a and from H-4 to C-3/C-1a/C-3a (Fig. 2), confirmed the identification of compound 4 as 1*H*-indole-3-carbaldehyde, previously reported from *Laminaria japonica* (Wang *et al.*, 2013).

The other isolates were identified as, 3-hydroxy-1-(4-hydroxy-3-methoxyphenyl)propan-1-one (3) (Jones *et al.*, 2000), *trans*-cinnamic acid (5) (Davidse *et al.*, 1990), and uracil (6) (Kitajima *et al.*, 1999), by comparison of their physical and spectral data with published values. Among the isolates, compound 1 was obtained as a major metabolite. To the best of our knowledge, compounds 2-6 were isolated for the first time from this plant source.

## 2. Evaluation of anti-inflammatory activity

The CHCl<sub>3</sub>-soluble portion from the aerial parts of *V. cinerea* exhibited potent inhibitory activity against TNF- $\alpha$ -induced NF- $\kappa$ B activity and NO production in LPS-stimulated RAW 264.7 cells with inhibition rates of 62.3 and 63.3%, respectively. Compounds 1-3 isolated from the CHCl<sub>3</sub> extract were also evaluated for their anti-inflammatory potential based on their ability to inhibit TNF- $\alpha$ -induced NF- $\kappa$ B activity and NO production, while compounds 4-6 could not be evaluated in this manner, due to their insufficient amounts available for testing. As shown in Table 1, when treated with a fixed concentration of 50  $\mu\text{M}$ , quercetin glycoside, 1 and phenyl propanoid, 3 exhibited moderate TNF- $\alpha$ -induced NF- $\kappa$ B inhibitory activity with IC<sub>50</sub> values of 7.5 and 11.5  $\mu\text{M}$ , respectively, compare to that of the positive control, tosyl phenylalanyl chloromethyl ketone (TPCK, IC<sub>50</sub>=3.8 M), whereas compounds 1-3 showed weak inhibitory activity on NO production.

**Table 1.** Inhibition effect of compounds 1 - 3 on the TNF- $\alpha$ -induced NF- $\kappa$ B activity and NO production in LPS-stimulated RAW 264.7 cells.

Compounds	Nitrite assay (NO)			NF- $\kappa$ B		
	%Inhibitory <sup>1)</sup>	%Survive <sup>2)</sup>	IC <sub>50</sub> (M)	%Inhibitory	%Survive	IC <sub>50</sub> (M)
Aerial parts	56.2 $\pm$ 1.5*	> 100	17.6 $\pm$ 0.8	35.5 $\pm$ 0.6*	> 100	nd
Roots	44.8 $\pm$ 2.5*	> 100	nd <sup>3)</sup>	8.7 $\pm$ 3.2*	> 100	nd
CH <sub>2</sub> Cl <sub>2</sub> partition	63.3 $\pm$ 3.7*	> 100	nd	62.3 $\pm$ 5.6*	> 100	nd
BuOH partition	55.4 $\pm$ 4.4*	> 100	nd	34.1 $\pm$ 1.5*	> 100	nd
1	23.8 $\pm$ 4.8	86.1 $\pm$ 5.7	nd	68.3 $\pm$ 8.8	80.0 $\pm$ 6.2	7.5 $\pm$ 2.5
2	15.1 $\pm$ 10.6	> 100	nd	61.0 $\pm$ 8.4**	94.6 $\pm$ 11.3	nd
3	26.7 $\pm$ 7.3**	> 100	nd	57.0 $\pm$ 6.6**	80.0 $\pm$ 8.6	11.5 $\pm$ 1.2
4	- <sup>4)</sup>	-	-	-	-	-
5	-	-	-	-	-	-
6	-	-	-	-	-	-
I-NMMA <sup>5)</sup>			25.1 $\pm$ 2.3			
TPCK <sup>6)</sup>						3.8 $\pm$ 0.6
BAY-11 <sup>6)</sup>						2.0 $\pm$ 0.3

1) % inhibition at 50  $\mu$ M, 2) % survival at concentration of 50  $\mu$ M, 3) nd; Not determined, 4) -; not tested, 5) positive control for NO, 6) positive control for NF- $\kappa$ B. Data represent means  $\pm$  SD of triplicate determinations from three separate experiments. Significantly different at \* $p$  < 0.05 and \*\* $p$  < 0.01 between groups.

Compound 3 has been reported to have weak activity on NO production (Min and Cuong, 2013), but inhibitory activity of 3 on TNF- $\alpha$ -induced NF- $\kappa$ B was evaluated first time in this study. In addition, quercetin and its derivatives have been found to possess NF- $\kappa$ B inhibitory and antioxidant effects (Kim *et al.*, 2013; Lee *et al.*, 2013).

In a previous research for cancer chemopreventive agents, major secondary metabolites, sesquiterpene lactones isolated from the flower of *V. cinerea* exhibited potent inhibitory effects on the tumor necrosis factor alpha (TNF- $\alpha$ )-induced NF- $\kappa$ B activity and lipopolysaccharide (LPS)-induced nitric oxide production using murine macrophage RAW 264.7 cells (Youn *et al.*, 2012). Although the sesquiterpene lactones have been reported to have a potential on the cancer chemopreventive activity in the previous study, the anti-inflammatory activity of several phenolic compounds isolated from the aerial parts of *V. cinerea* can be attributed, at least in part, to inhibition of TNF- $\alpha$ -induced NF- $\kappa$ B activity.

#### ACKNOWLEDGMENTS

This study was supported by a grant to the Korea Polar Research Institute (KOPRI), under a project PE16350. We

thank H. S. Shin, National center for inter University Research Facilities, Seoul National University, for the provision of the Mass Spectrometry Facility used in this study.

#### REFERENCES

- Anggård E. (1994). Nitric oxide: Mediator, murderer, and medicine. *Lancet*. 343:1199-1206.
- Baldwin AS. (2001). Control of oncogenesis and cancer therapy resistance by the transcription factor NF- $\kappa$ B. *Journal of Clinical Investigation*. 107:241-246.
- Beeran AA, Maliyakkal N, Rao CM and Udupa N. (2014). The enriched fraction of *Vernonia cinerea* L. induces apoptosis and inhibits multi-drug resistance transporters in human epithelial cancer cells. *Journal of Ethnopharmacology*. 158:33-42.
- Balasundram N, Sundram K and Samman S. (2006). Phenolic compounds in plants and agri-industrial by-products: Antioxidant activity, occurrence, and potential uses. *Food Chemistry*. 99:191-203.
- Chea A, Hout S, Long C, Marcourt L, Faure R, Azas N and Elias R. (2006). Antimalarial activity of sesquiterpene lactones from *Vernonia cinerea*. *Chemical and Pharmaceutical Bulletin*. 54:1437-1439.
- Choi YH, Kim JH, Kim MJ, Han SS and Rim YS. (2000). Antioxidative compounds in leaves of *Castanea crenata* S. et Z. *Korean Journal of Medicinal Crop Sciences*. 8:373-377.
- Dastur JF. (1977). Medicinal plants of India and Pakistan. D.B Taraporevala Sons and Co., Ltd. Maharashtra, India. p.174.
- Davide PA, Dillen JLM, Heyns AM, Modro TA and Rooyen PHV. (1990). Photochromic systems. Part 1. Structural and spectroscopic study of photochromically active products of Stobbe condensation.

- 2,3-Dibenzylidenesuccinic acid and its anhydride. *Canadian Journal of Chemistry*. 68:741-746.
- Han JT, Bang MH, Chun OK, Kim DO, Lee CY and Baek NI.** (2004). Flavonol glycosides from the aerial parts of *Aceriphyllum rossii* and their antioxidant activities. *Archives of Pharmacal Research*. 27:390-395.
- Jones L, Bartholomew B, Latif Z, Sarker SD and Nash RJ.** (2000). Constituents of *Cassia laevigata*. *Fitoterapia*. 71:580-583.
- Kim BH, Choi JS, Yi EH, Lee JK, Won C, Ye SK and Kim MH.** (2013). Relative antioxidant activities of quercetin and its structurally related substances and their effects on NF- $\kappa$ B/CRE/AP-1 signaling in murine macrophages. *Molecules and Cells*. 35:410-420.
- Kirtikar KR, Basu BD and an I.C.S.** (1975). Indian medicinal plants. Bishen Singh Mahendra Pal Singh. Dehradun, India. p.1322.
- Kitajima J, Ishikawa T, Tanaka T and Ida Y.** (1999). Water-soluble constituents of fennel. IX. Glucides and nucleosides. *Chemical and Pharmaceutical Bulletin*. 47:988-992.
- Kondratyuk TP, Park EJ, Yu R, van Breemen RB, Asolkar RN, Murphy BT, Fenical W and Pezzuto JM.** (2012). Novel marine phenazines as potential cancer chemopreventive and anti-inflammatory agents. *Marine Drugs*. 10:451-464.
- Kone WM and Kande B.** (2012). Qualitative analysis of the pyrrolizidine alkaloids from 11 Asteraceae and Boraginaceae used in traditional medicine in Cote d'Ivoire. *Research Journal of Phytochemistry*. 6:75-83.
- Kuo YH, Kuo YJ, Yu AS, Wu MD, Ong CW, Kuo LMY, Huang JT, Chen CF and Li SY.** (2003). Two novel sesquiterpene lactones, cytotoxic vernolide-A and -B, from *Vernonia cinerea*. *Chemical and Pharmaceutical Bulletin*. 51:425-426.
- Lee CS, Jeong EB, Kim YJ, Lee MS, Seo SJ, Park KH and Lee MW.** (2013). Quercetin-3-O-(2"-galloyl)- $\alpha$ -L-rhamnopyranoside inhibits TNF- $\alpha$ -activated NF- $\kappa$ B-induced inflammatory mediator production by suppressing ERK activation. *International Immunopharmacology*. 16:481-487.
- Manach C, Mazur A and Scalbert A.** (2005). Polyphenols and prevention of cardiovascular diseases. *Current Opinion in Lipidology*. 16:77-84.
- Masuda T and Jitoe A.** (1995). Phenylbutenoid monomers from the rhizomes of *Zingiber cassumunar*. *Phytochemistry*. 39:459-461.
- Middleton E Jr, Kandaswami C and Theoharis TC.** (2000). The effects of plant flavonoids on mammalian cells: Implications for inflammation, heart disease, and cancer. *Pharmacological Reviews*. 52:673-751.
- Min BS and Cuong TD.** (2013). Phenolic compounds from *Caesalpinia sappan* and their inhibitory effects on LPS-induced NO production in RAW 264.7 cells. *Natural Product Sciences*. 19:201-205.
- Misra TN, Singh RS, Upadhyay J and Srivastava R.** (1984a). Chemical constituents of *Vernonia cinerea*, part I. Isolation and spectral studies of triterpenes. *Journal of Natural Products*. 47:368-372.
- Misra TN, Singh RS, Upadhyay J and Srivastava R.** (1984b). Isolation of a natural sterol and an aliphatic acid from *Vernonia cinerea*. *Phytochemistry*. 23:415-417.
- Moncada SRMJ, Palmer RML and Higgs EA.** (1991). Nitric oxide: Physiology, pathophysiology, and pharmacology. *Pharmacological Reviews*. 43:109-142.
- Park EJ, Kondratyuk TP, Morrell A, Kiselev E, Conda-Sheridan M, Cushman M, Ahn S, Choi Y, White JJ, van Breemen RB and Pezzuto JM.** (2011). Induction of retinoid X receptor activity and consequent upregulation of p21<sup>WAF1/CIP1</sup> by indenoisoquinolines in MCF7 cells. *Cancer Prevention Research*. 4:592-607.
- Pratheeshkumar P and Kuttan G.** (2011). Effect of vernolide-A, a sesquiterpene lactone from *Vernonia cinerea* L., on cell-mediated immune response in B16F-10 metastatic melanoma-bearing mice. *Immunopharmacology and Immunotoxicology*. 33:533-538.
- Tandon M, Shukla YN, Tripathi AK and Singh SC.** (1998). Insect antifeedant principles from *Vernonia cinerea*. *Phytotherapy Research*. 12:195-199.
- Wang C, Yang Y, Mei Z and Yang X.** (2013). Cytotoxic compounds from *Laminaria japonica*. *Chemistry of Natural Compounds*. 49:699-701.
- Youn UJ, Park EJ, Kondratyuk TP, Simmons CJ, Borris RP, Tanamatayarat P, Wongwiwatthanakut S, Toyama O, Songsak T, Pezzuto JM and Chang LC.** (2012). Anti-inflammatory sesquiterpene lactones from the flower of *Vernonia cinerea*. *Bioorganic and Medicinal Chemistry Letters*. 22:5559-5562.